

SPECIFICATION

Glycerol

for GC

REQUIREMENT	SPECIFICATION
Assay (HOCH ₂ CHOHCH ₂ OH) (GC)	min.99.5%(area/area)
Appearance (color)	max.10Hazen units
Water	max.1.0%(mass/mass)
Residue after ignition (as sulfate)	max.0.005%(mass/mass)
Acidity (as CH ₃ COOH)	max.0.002%(mass/mass)
Alkalinity (as NH ₃)	max.4ppm(mass/mass)
Chloride (Cl)	max.1ppm(mass/mass)
Sulfate (SO ₄)	max.5ppm(mass/mass)
Lead (Pb)	max.1ppm(mass/mass)
Iron (Fe)	max.1ppm(mass/mass)
Arsenic (As)	max.1ppm(mass/mass)
Ammonium (NH ₄)	max.5ppm(mass/mass)
Acrolein and sugars	to pass test
Fatty acid esters (as glycerol tributryate)	max.0.05%(mass/mass)
Aldehyde (as HCHO)	max.10ppm(mass/mass)
Substances darkened by sulfuric acid	to pass test
Ethylene glycol (HOCH ₂ CH ₂ OH)	max.0.02%(mass/mass)
Diethylene glycol [(HOC ₂ H ₄) ₂ O]	max.0.02%(mass/mass)

JIS K 8295:2020