

(90 × 210mm Size)

FUJIFILM

Wako

Code No. 180-02991 (100 μ L)
186-02993 (100 μ L × 5)

rBC2LCN-FITC (AiLecS1-FITC)

rBC2LCN is a recombinant protein of N-terminal domain of BC2L-C and has been reported as a marker of undifferentiated human pluripotent stem cells (hPSCs), human ES cells and human iPSC cells. This product is a FITC-labeled rBC2LCN.

rBC2LCN specifically recognize a sugar chain which exists on the surface of hPSCs. When you add rBC2LCN-FITC to the medium which is used for hPSCs culture, you can stain the living hPSCs. rBC2LCN-FITC is not toxic to cells, therefore you can culture hPSCs without any effect on the growth and pluripotency in the medium containing rBC2LCN-FITC.

This product is for laboratory research use only ; use in any such application is the responsibility of the user.

Formulation : Phosphate Buffered Saline sterilized with 0.1 μ m filter

Working Dilution : Live Cell Imaging 1:100~1:1,000
Flow Cytometry 1:100~1:1,000

Protocol for Live Cell Imaging

- 1) Culture hPSCs in appropriate culture conditions.
- 2) Add 1~10 μ L of rBC2LCN-FITC per medium 1 mL.
- 3) Incubate the cells over 30 min under appropriate conditions.
- 4) Replace the medium containing rBC2LCN-FITC to a fresh medium or D-PBS(-).
- 5) View and analyze the cells on an imaging instrument (excitation 495 nm, emission 520 nm).
※ Step 4 may be omitted.

Protocol for Flow Cytometry

- 1) Dissociate the hPSCs and wash with FCM* buffer once. Centrifuge and discard the supernatant.
- 2) Resuspend the cells with FCM* buffer to prepare cell suspension of approximately 5×10^6 cells/mL.
- 3) Add 1-10 μ L of rBC2LCN-FITC per 1 mL of cell suspension.
- 4) Incubate the cells in the dark over 30 min at room temperature.
- 5) Centrifuge the cell suspension at 1,000 rpm for 3 min, and discard the supernatant.
- 6) Wash the cells with 1 mL of FCM* buffer, centrifuge the cell suspension at 1,000 rpm for 3 min and discard the supernatant.
- 7) Suspend the washed cells in appropriate volume of FCM* buffer.
- 8) Analyze the cells by flow cytometry.
* FCM buffer : D-PBS(-) containing 10 mmol/L EDTA and 1% FBS, and so on.

- 1/2 -

Attention :

- 1) Staining with the rBC2LCN-FITC persists for a few days even after removing the rBC2LCN-FITC.
- 2) For continuous observations, it might be better to add the same concentration of rBC2LCN-FITC at the daily medium change. Cytotoxicity of the rBC2LCN-FITC has not been observed.
- 3) rBC2LCN-FITC is applicable to not only live cell imaging, but also staining fixed cells.
- 4) The signal of background might become higher when the cells are cultured with the medium containing serum.

Storage : Store at -20 °C, in the dark
After thawing, store at 2~10 °C. Avoid repeating freeze-thaw.

Package :

Code No.	Packaging
180-02991	100 μ L
186-02993	100 μ L × 5

FUJIFILM Wako Pure Chemical Corporation

1-2, Doshomachi 3-Chome, Chuo-Ku, Osaka 540-8605, Japan
Telephone : + 81-6-6203-3741
Facsimile : + 81-6-6201-5964
<http://www.wako-chem.co.jp>

FUJIFILM Wako Chemicals U.S.A. Corporation

1600 Bellwood Road
Richmond, VA 23237
U.S.A.
Telephone : + 1-804-271-7677
Facsimile : + 1-804-271-7791
<http://www.wakousa.com>

FUJIFILM Wako Chemicals Europe GmbH

Fuggerstrasse 12
D-41468 Neuss
Germany
Telephone : + 49-2131-3111-0
Facsimile : + 49-2131-311100
<http://www.wako-chemicals.de>

1801KA1

- 2/2 -