

## SPECIFICATION

## Potassium Chloride

for Measurement of Electric Conductivity

REQUIREMENT	SPECIFICATION
* Appearance	White, crystals ~ crystalline powder
Assay (KCl) (after drying at 500°C)	min.99.9%(mass/mass)
Solubility in water	to pass test
Loss on drying at 110°C	max.0.1%(mass/mass)
pH of a 50g/L solution at 25°C	5.0~8.0
Bromide (Br)	max.0.01%(mass/mass)
Iodide (I)	max.0.002%(mass/mass)
Phosphate (PO <sub>4</sub> )	max.5ppm(mass/mass)
Sulfate (SO <sub>4</sub> )	max.0.002%(mass/mass)
Nitrogen compounds (as N)	max.0.001%(mass/mass)
Sodium (Na)	max.0.02%(mass/mass)
Copper (Cu)	max.2ppm(mass/mass)
Magnesium (Mg)	max.0.002%(mass/mass)
Calcium (Ca)	max.0.002%(mass/mass)
Barium (Ba)	max.0.001%(mass/mass)
Lead (Pb)	max.2ppm(mass/mass)
Iron (Fe)	max.2ppm(mass/mass)

JIS K 8121:2023(\*:Additional test performed by Wako but not required by JIS)